



BULLETIN OF MATHEMATICS AND STATISTICS RESEARCH

A Peer Reviewed International Research Journal



GENERALIZED HYERS-ULAM -RASSISA TYPE STABILITY OF AN ADDITIVE(Γ_1 , Γ_2) - FUNCTIONAL INEQUALITIES WITH THREE VARIABLES IN COMPLEX BANACH SPACE

LY VAN AN

Faculty of Mathematics Teacher Education, Tay Ninh University, Vietnam.

Email: lyvanan145@gmail.com(lyvananvietnam).

DOI:[10.33329/bomsr.10.4.8](https://doi.org/10.33329/bomsr.10.4.8)



ABSTRACT

In this paper we study to solve the of additive (Γ_1, Γ_2) -functional inequality with three variables and their Hyers-Ulam stability. First are investigated in complex Banach spaces with a fixed point method and last are investigated in complex Banach spaces with a direct method. These are the main results of this paper.

LY VAN AN

Keywords: Additive (Γ_1, Γ_2) -Functional inequality; Fixed Point Method; Direct Method; Banach Space; Hyers Ulam Stability.

Mathematics Subject Classification : 46S10, 39B62, 39B52, 47H10,

Let \mathbb{X} and \mathbb{Y} be a normed spaces on the same field \mathbb{K} , and $f: \mathbb{X} \rightarrow \mathbb{Y}$. We use the notation $\| \cdot \|$ for all the norm on both \mathbb{X} and \mathbb{Y} . In this paper, we investigate additive (Γ_1, Γ_2) -functional inequality when \mathbb{X} be a real or complex normed space and \mathbb{Y} a complex Banach spaces. We solve and prove the Hyers-Ulam stability of following additive (Γ_1, Γ_2) -functional inequality.

$$\begin{aligned}
 & \left\| f\left(\frac{x+y}{2}+z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\
 & \leq \left\| \Gamma_1\left(f\left(\frac{x+y}{2}+z\right) - f\left(\frac{x+y}{2}-z\right) - 2f\left(\frac{x+y}{2}\right)\right) \right\|_{\mathbb{Y}} \\
 & + \left\| \Gamma_2\left(2f\left(\frac{x+y}{2^2}+\frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) \right. \right. \\
 & \quad \left. \left. - f(z)\right) \right\|_{\mathbb{Y}}
 \end{aligned} \tag{1.1}$$

In which Γ_1, Γ_2 are fixed non zero complex numbers with $G(\Gamma_1, \Gamma_2)$ -functional inequality.

Note that in the preliminaries we just recap some of the most essential properties for the above problem and for the specific problem, please see the document. The Hyers- Ulam stability was first investigated for functional equation of Ulam in [28] concerning the stability of group homomorphisms.

The functional equation

$$f(x + y) = f(x) + f(y)$$

is called the Cauchy equation. In particular, every solution of the Cauchy equation is said to be an additive mapping.

The Hyers [13] gave first affirmative partial answer to the equation of Ulam in Banach spaces. After that, Hyers' Theorem was generalized by Aoki[1] additive mappings and by Rassias [26] for linear mappings considering an unbounded Cauchy difference. A generalization of the Rassias theorem was obtained by Găvruta [10] by replacing the unbounded Cauchy difference by a general control function in the spirit of Rassias' approach.

The stability of quadratic functional equation was proved by Skof [27] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Park [24], [25] defined additive γ -functional inequalities and proved the Hyers-Ulam stability of the additive γ -functional inequalities in Banach spaces and non-Archimedean Banach spaces. The stability problems of various functional equations have been extensively investigated by a number of authors on the world. We recall a fundamental result in fixed point theory.

Recently, in [3],[4],[21],[22],[24],[25] the authors studied the Hyers-Ulam stability for the following functional inequalities

$$\left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\| \leq \left\| f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - \frac{1}{2}f\left(\frac{x+y}{2}\right) - f(z) \right\| \quad (1.2)$$

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - \frac{1}{2}f\left(\frac{x+y}{2}\right) - \frac{1}{2}f(z) \right\| \\ & \leq \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\| \end{aligned} \quad (1.3)$$

$$\|f(x+y) - f(x) - f(y)\| \leq \left\| \rho(2f\left(\frac{x+y}{2}\right) - f(x) - f(y)) \right\| \quad (1.4)$$

$$\left\| 2f\left(\frac{x+y}{2}\right) - f(x) - f(y) \right\| \leq \|\rho(f(x+y) - f(x) - f(y))\| \quad (1.5)$$

and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - f(z) - 2f\left(\frac{x+y}{2}\right) - 2f(z) \right\| \\ & \leq \left\| \beta(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - 2f\left(\frac{x+y}{2^2} - \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) \right. \\ & \quad \left. - f(z)) \right\| \end{aligned} \quad (1.6)$$

$$\begin{aligned} & \left\| 2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) + 2f\left(\frac{x+y}{2^2} - \frac{z}{2}\right) - f(z) - f\left(\frac{x+y}{2}\right) - 2f(z) \right\| \\ & \leq \left\| \beta(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right. \\ & \quad \left. - 2f(z)) \right\| \end{aligned} \quad (1.7)$$

Finaly

$$\begin{aligned} &\leq \|\beta_1(f(x+y) + f(x-y) - 2f(x))\| \\ &+ \left\| \beta_2(2f\left(\frac{x+y}{2}\right) - f(x) - f(y)) \right\| \quad (1.8) \end{aligned}$$

in complex Banach spaces

In this paper, we solv e and proved the Hyers-Ulam stability for (Γ_1, Γ_2) - functional inequalities (1.1), i.e., the (Γ_1, Γ_2) - functional inequalities with three variables. Under suitable assumptions on spaces \mathbb{X} and \mathbb{Y} , we will prove that the mappings satisfying the (Γ_1, Γ_2) - functional inequati lies (1. 1). Thus, the results in this paper are generalization of those in [3],[4],[14],[21] for (Γ_1, Γ_2) - functional inequati lies with three variables.

The paper is organized as follows: In section preliminarier we remind some basic nota- tions in [3],[7] such as complete generalized metric space and Solutions of the inequalities.

Section 3: In this section, I use the method of the fixed to prove the Hyers-Ulam stability of the additive (Γ_1, Γ_2) -functional inequalities (1.1) when \mathbb{X} be a real or complete normed space and \mathbb{Y} complex Banach space.

Section 4: In this section, I use the method of directly determining the solution for (1.1) when \mathbb{X} be a real or complete normed space and \mathbb{Y} complex Banach space.

2.1. complete generalized metric space and Solutions of the inequalities.

Theorem 2.1. Let (X, d) be a complete generalized metric space and let $J : X \rightarrow X$ be a strictly contractive mapping with Lipschitz constant $L < 1$. Then for each given element $x \in X$, either

$$d(J^n, J^{n+1}) = \infty$$

for all nonnegative integers n or there exists a positive integer n_0 such that

- (1) $d(J^n, J^{n+1}) < \infty, \forall n \geq n_0$;
- (2) The sequence $\{J^n x\}$ converges to a fixed point y^* of J ;
- (3) y^* is the unique fixed point of J in the set
- (4) $Y = \{y \in X \mid d(J^n, J^{n+1}) < \infty\}$;
- (5) $d(y, y^*) \leq \frac{1}{1-l} d(y, J_y) \forall y \in Y$

Solutions of the inequalities. The functional equation

$$f(x + y) = f(x) + f(y)$$

is called the cauchuy equation. In particular, every solution of the cauchuy equation is said to be an additive mapping.

3. ESTABLISH THE SOLUTION OF THE ADDITIVE (Γ_1, Γ_2) -FUNCTION INEQUALITIES USING A FIXED POINT METHOD

Now, we first study the solutions of (1.1). Note that for these inequalities, when \mathbb{X} be a real or complete normed space and \mathbb{Y} complex Banach space.

Lemma 3.1. A mapping $f: \mathbb{X} \rightarrow \mathbb{Y}$ satisfies $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right)) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) \right. \\ & \left. - f(z)) \right\|_{\mathbb{Y}} \end{aligned} \quad (3.1)$$

for all $x, y, z \in X$ if and only if $f: X \rightarrow Y$ is additive

Proof. Assume that $f: X \rightarrow Y$ satisfies (3.1)

Letting $x = y = z$ in (3.1), we get

$$\begin{aligned} \|f(2x) - 2f(x)\|_{\mathbb{Y}} \\ \leq |\Gamma_1| \|f(2x) - 2f(x)\|_{\mathbb{Y}} \end{aligned}$$

and so $f(2x) = 2f(x)$ for all $x \in X$.

Thus

$$f(x/2) = 1/2f(x) \quad (3.2)$$

for all $x \in X$ It follows from (3.1) and (3.2) that

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1\left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right)\right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z)) \right\|_{\mathbb{Y}} \\ & = \left\| \Gamma_1\left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right)\right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2(f\left(\frac{x+y}{2}\right) + z) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \end{aligned} \quad (3.3)$$

and so

$$\begin{aligned} & (1 - |\Gamma_2|) \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq |\Gamma_1| \left\| \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right)\right) \right\|_{\mathbb{Y}} \end{aligned} \quad (3.4)$$

Letting $u = \frac{x+y}{2} + z$; $v = \frac{x+y}{2} - z$ in (3.4), we get

$$\begin{aligned} & (1 - |\Gamma_2|) \left\| f(u) - f\left(\frac{u+v}{2}\right) - f\left(\frac{u-v}{2}\right) \right\|_{\mathbb{Y}} \\ & \leq |\Gamma_1| \left\| f(u) - f(v) - 2f\left(\frac{u+v}{2}\right) \right\|_{\mathbb{Y}} \end{aligned} \quad (3.5)$$

for all $u, v \in X$ and so

$$\begin{aligned} & \frac{1}{2}(1 - |\Gamma_2|) \|f(u + v) - f(u - v) + 2f(u)\|_{\mathbb{Y}} \\ & \leq |\Gamma_1| \|f(u + v) - f(u) - f(v)\|_{\mathbb{Y}} \end{aligned} \quad (3.6)$$

for all $u, v \in X$ it follows from (3.4) and (3.5), that

$$\begin{aligned} & \frac{1}{2}(1 - |\Gamma_2|)^2 \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq |\Gamma_1|^2 \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \end{aligned} \quad (3.7)$$

Since $\sqrt{2} |\beta_1| + |\beta_2| < 1$ and so

$$f\left(\frac{x+y}{2} + z\right) = f\left(\frac{x+y}{2}\right) + f(z)$$

for all $x, y, z \in X$. Thus f is additive

Theorem 3.2. Let $\varphi : \mathbb{X}^3 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right) \leq \frac{L}{2} \varphi(x, y, z) \quad (3.8)$$

for all $x, y, z \in \mathbb{X}$. Let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \varphi(x, y, z) \end{aligned} \quad (3.9)$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique mapping $h : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\|f(x) - h(x)\|_{\mathbb{Y}} \leq \frac{L}{2(1-L)(1-|\Gamma_1|)} \varphi(x, x, x) \quad (3.10)$$

for all $x \in \mathbb{X}$

Proof. Letting $x = y = z$ in (3.9), we get

$$(1 - |\Gamma_1|) \left\| f(2x) - 2f(x) \right\|_{\mathbb{Y}} \leq \varphi(x, x, x) \quad (3.11)$$

for all $x \in \mathbb{X}$.

Consider the set

$$\mathbb{S} := \left\{ h : \mathbb{X} \rightarrow \mathbb{Y}, h(0) = 0 \right\}$$

and introduce the generalized metric on \mathbb{S} :

$$d(g, h) := \inf \left\{ \lambda \in \mathbb{R} : \|g(x) - h(x)\| \leq \lambda \varphi(x, x, x), \forall x \in X \right\},$$

where, as usual, $\inf \phi = +\infty$. It is easy to show that (\mathbb{S}, d) is complete (see [16]). Now we consider the linear mapping $J : \mathbb{S} \rightarrow \mathbb{S}$ such that

$$Jg(x) := 2g\left(\frac{x}{2}\right)$$

for all $x \in \mathbb{X}$. Let $g, h \in \mathbb{S}$ be given such that $d(g, h) = \epsilon$ then

$$\left\| g(x) - h(x) \right\| \leq \epsilon \varphi(x, x, x)$$

for all $g, h \in \mathbb{S}$. It follows from (3.11) that

$$\left\| f(x) - 2f\left(\frac{x}{2}\right) \right\| \leq \frac{1}{1 - |\beta_1|} \varphi\left(\frac{x}{2}, \frac{x}{2}, \frac{x}{2}\right) \leq \frac{L}{2(1 - |\Gamma_1|)} \varphi(x, x, x)$$

for all $x \in \mathbb{X}$. So $d(f, Jf) \leq \frac{L}{2(1 - |\Gamma_1|)}$ for all $x \in \mathbb{X}$. By Theorem 2.3, there exists a mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ satisfying the following:

- (1) H is a fixed point of J , ie.,

$$H\left(\frac{x}{2}\right) = 2H\left(\frac{x}{2}\right) \quad (3.12)$$

for all $x \in \mathbb{X}$. The mapping H is a unique fixed point J in the set

$$\mathbb{M} = \left\{ g \in \mathbb{S} : d(f, g) < \infty \right\}$$

This implies that H is a unique mapping satisfying (3.12) such that there exists a $\lambda \in (0, \infty)$ satisfying

$$\left\| f(x) - H(x) \right\| \leq \lambda \varphi(x, x, x)$$

(2) for all $x \in \mathbb{X}$
 $d(J^l f, H) \rightarrow 0$ as $l \rightarrow \infty$. This implies equality

$$\lim_{l \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right) = H(x)$$

(3) for all $x \in \mathbb{X}$
 $d(f, H) \leq \frac{1}{1-L} d(f, Jf)$. which implies

$$\left\| f(x) - H(x) \right\| \leq \frac{L}{2(1-L)(1-|\Gamma_1|)} \varphi(x, x, x)$$

for all $x \in X$. It follows (3.8) and (3.9) that

$$\begin{aligned} & \left\| H\left(\frac{x+y}{2} + z\right) - H\left(\frac{x+y}{2}\right) - H(z) \right\| \\ &= \lim_{n \rightarrow \infty} 2^n \left\| f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) - f\left(\frac{x+y}{2^{n+1}}\right) - f\left(\frac{z}{2^n}\right) \right\| \\ &\leq \lim_{n \rightarrow \infty} 2^n |\Gamma_1| \left\| f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) + f\left(\frac{x+y}{2^{n+1}} - \frac{z}{2^n}\right) - 2f\left(\frac{x+y}{2^{n+1}}\right) \right\| \\ &+ \lim_{n \rightarrow \infty} 2^n |\Gamma_2| \left\| 2f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) + f\left(\frac{x+y}{2^n}\right) - f\left(\frac{z}{2^n}\right) \right\| + \lim_{n \rightarrow \infty} 2^n \varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right) \\ &= \left\| \Gamma_1 \left(H\left(\frac{x+y}{2} + z\right) + H\left(\frac{x+y}{2} - z\right) - 2H\left(\frac{x+y}{2}\right) \right) \right\| + \left\| \Gamma_2 \left(2H\left(\frac{x+y}{2^2} + \frac{z}{2}\right) \right. \right. \\ &\quad \left. \left. - H\left(\frac{x+y}{2}\right) - H(z) \right) \right\|_{\mathbb{Y}} \end{aligned}$$

for all $x, y, z \in X$. So

$$\begin{aligned} & \left\| H\left(\frac{x+y}{2} + z\right) - H\left(\frac{x+y}{2}\right) - H(z) \right\| \\ &\leq \left\| \Gamma_1 \left(H\left(\frac{x+y}{2} + z\right) + H\left(\frac{x+y}{2} - z\right) - 2H\left(\frac{x+y}{2}\right) \right) \right\| + \left\| \Gamma_2 \left(2H\left(\frac{x+y}{2^2} + \frac{z}{2}\right) \right. \right. \\ &\quad \left. \left. - H\left(\frac{x+y}{2}\right) - H(z) \right) \right\|_{\mathbb{Y}} \end{aligned}$$

$$H\left(\frac{x+y}{2}\right) + H(z) = H\left(\frac{x+y}{2} + z\right)$$

for all $x, y, z \in X$. By Lemma 3.1, the mapping $H : X \rightarrow Y$ is additive.

Theorem 3.3. Let $\varphi : \mathbb{X}^3 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y, z) \leq \frac{L}{2}\varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right) \quad (3.13)$$

for all $x, y, z \in \mathbb{X}$. Let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \varphi(x, y, z) \end{aligned} \quad (3.14)$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique mapping $h : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\left\| f(x) - h(x) \right\|_{\mathbb{Y}} \leq \frac{L}{2(1-L)(1-|\Gamma_1|)} \varphi(x, x, x) \quad (3.15)$$

for all $x \in \mathbb{X}$

Proof. Letting $x = y = z$ in (3.14), we get

$$(1 - |\Gamma_1|) \left\| f(2x) - 2f(x) \right\|_{\mathbb{Y}} \leq \varphi(x, x, x) \quad (3.16)$$

and so

$$\left\| f(x) - \frac{1}{2}f(2x) \right\|_{\mathbb{Y}} \leq \frac{1}{2(1 - |\Gamma_1|)} \varphi(x, x, x) \quad (3.17)$$

for all $x \in \mathbb{X}$.

Let (S, d) be the generalized metric space defined in the proof of Theorem 3.2
Now we consider the mapping $J : S \rightarrow S$ such that

$$Jg(x) := \frac{1}{2}g(2x)$$

for all $x \in X$.

It follows (3.17) that $d(f, Jf) \leq \frac{1}{2(1 - |\Gamma_1|)}$. So

$$\left\| f(x) - H(x) \right\| \leq \frac{1}{2(1 - L)(1 - |\Gamma_1|)} \varphi(x, x, x) \quad (3.18)$$

for all $x \in X$.

The rest of the proof is similar to proof of Theorem 3.2.

From proving the theorems we have consequences:

Corollary 3.4. Let $r > 1$ and θ be nonnegative real numbers and let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \theta \left(\|x\|^r + \|y\|^r + \|z\|^r \right) \end{aligned} \quad (3.19)$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbb{Y}} \leq \frac{3\theta}{(2 - 2^r)(1 - |\Gamma_1|)} \varphi(x, x, x) \quad (3.20)$$

for all $x \in \mathbb{X}$

Corollary 3.5. Let $r < 1$ and θ be nonnegative real numbers and let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and

$$\left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}}$$

$$\begin{aligned}
&\leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\
&+ \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \theta \left(\|x\|^r + \|y\|^r + \|z\|^r \right)
\end{aligned} \tag{3.21}$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\|f(x) - H(x)\|_{\mathbb{Y}} \leq \frac{3\theta}{(2-2^r)(1-|\Gamma_1|)} \varphi(x, x, x) \tag{3.22}$$

for all $x \in \mathbb{X}$

4. ESTABLISH THE SOLUTION OF THE ADDITIVE (Γ_1, Γ_2) -FUNCTION INEQUALITIES USING A DIRECT METHOD

Next, we study the solutions of (1.1). Note that for these inequalities, when X be a real or complete normed space and Y complex Banach space.

Theorem 4.1. Let $\varphi : X^3 \rightarrow [0, \infty)$ be a function and let $f : X \rightarrow Y$ be a mapping such that

$$\psi(x, y, z) := \sum_{j=1}^{\infty} 2^j \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}, \frac{z}{2^j}\right) < \infty \tag{4.1}$$

and let $f : X \rightarrow Y$ be a mapping $f(0) = 0$ and

$$\begin{aligned}
&\left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\
&\leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\
&+ \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \varphi(x, y, z)
\end{aligned} \tag{4.2}$$

for all $x, y, z \in \mathbb{X}$.
Then there exists a unique additive mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\|f(x) - H(x)\|_{\mathbb{Y}} \leq \frac{1}{2(1-|\Gamma_1|)} \psi(x, x, x) \tag{4.3}$$

for all $x \in \mathbb{X}$

Proof. Let $x = y = z$ in (4.2), we get

$$(1-|\Gamma_1|) \|f(2x) - 2f(x)\| \leq \varphi(x, x, x) \tag{4.4}$$

for all $x \in X$. So

$$\left\| f(x) - 2f\left(\frac{x}{2}\right) \right\| \leq \frac{1}{1 - |\Gamma_1|} \varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right)$$

for all $x \in X$. Hence

$$\begin{aligned} & \left\| 2^l f\left(\frac{x}{2^l}\right) - 2^m f\left(\frac{x}{2^m}\right) \right\| \\ & \leq \sum_{j=l}^{m-1} \left\| 2^j f\left(\frac{x}{2^j}\right) - 2^{j+1} f\left(\frac{x}{2^{j+1}}\right) \right\| \\ & \leq \sum_{j=l}^{m-1} \frac{2^j}{2(1 - |\Gamma_1|)} \varphi\left(\frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}, \frac{x}{2^{j+1}}\right) \end{aligned} \quad (4.5)$$

for all nonnegative integers m and l with $m > l$ and all $x \in \mathbb{X}$. It follows from (4.5) that the sequence $\left\{ 2^n f\left(\frac{x}{2^n}\right) \right\}$ is a Cauchy sequence for all $x \in \mathbb{X}$. Since \mathbb{Y} is complete, the sequence $\left\{ 2^n f\left(\frac{x}{2^n}\right) \right\}$ converges. So one can define the mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ by

$$H(x) := \lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x)$$

It follows from (4.1) and (4.2) that

$$\begin{aligned} & \left\| H\left(\frac{x+y}{2} + z\right) - H\left(\frac{x+y}{2}\right) - H(z) \right\| \\ & = \lim_{n \rightarrow \infty} 2^n \left\| f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) - f\left(\frac{x+y}{2^{n+1}}\right) - f\left(\frac{z}{2^n}\right) \right\| \\ & \leq \lim_{n \rightarrow \infty} 2^n |\Gamma_1| \left\| f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) + f\left(\frac{x+y}{2^{n+1}} - \frac{z}{2^n}\right) - 2f\left(\frac{x+y}{2^{n+1}}\right) \right\| \\ & \quad + \lim_{n \rightarrow \infty} 2^n |\Gamma_2| \left\| 2f\left(\frac{x+y}{2^{n+1}} + \frac{z}{2^n}\right) + f\left(\frac{x+y}{2^n}\right) - f\left(\frac{z}{2^n}\right) \right\| + \lim_{n \rightarrow \infty} 2^n \varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right) \\ & = \left\| \Gamma_1 \left(H\left(\frac{x+y}{2} + z\right) + H\left(\frac{x+y}{2} - z\right) - 2H\left(\frac{x+y}{2}\right) \right) \right\| + \left\| \Gamma_2 \left(2H\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - H\left(\frac{x+y}{2}\right) - H(z) \right) \right\|_{\mathbb{Y}} \end{aligned}$$

for all $x, y, z \in X$. So

$$\begin{aligned} & \left\| H\left(\frac{x+y}{2} + z\right) - H\left(\frac{x+y}{2}\right) - H(z) \right\| \\ & \leq \left\| \Gamma_1\left(H\left(\frac{x+y}{2} + z\right) + H\left(\frac{x+y}{2} - z\right) - 2H\left(\frac{x+y}{2}\right)\right) \right\| + \left\| \Gamma_2\left(2H\left(\frac{x+y}{2^2} + \frac{z}{2}\right) \right. \right. \\ & \quad \left. \left. - H\left(\frac{x+y}{2}\right) - H(z)\right) \right\|_{\mathbb{Y}} \\ & \quad H\left(\frac{x+y}{2}\right) + H(z) = H\left(\frac{x+y}{2} + z\right) \end{aligned}$$

for all $x, y, z \in X$. By Lemma 3.1, the mapping $H : X \rightarrow Y$ is additive.

Now, let $T : \mathbb{X} \rightarrow \mathbb{Y}$ be another additive mapping satisfying (4.3). Then we have

$$\begin{aligned} \left\| H(x) - T(x) \right\| &= \left\| 2^q H\left(\frac{x}{2^q}\right) - 2^q T\left(\frac{x}{2^q}\right) \right\| \\ &\leq \left\| 2^q H\left(\frac{x}{2^q}\right) - 2^q f\left(\frac{x}{2^q}\right) \right\| + \left\| 2^q T\left(\frac{x}{2^q}\right) - 2^q f\left(\frac{x}{2^q}\right) \right\| \\ &\leq \frac{2^q}{1 - |\Gamma_1|} \psi\left(\frac{x}{2^q}, \frac{x}{2^q}, \frac{x}{2^q}\right) \end{aligned}$$

which tends to zero as $q \rightarrow \infty$ for all $x \in \mathbb{X}$. So we can conclude that $H(x) = T(x)$ for all $x \in X$. This proves the uniqueness of H .

Theorem 4.2. Let $\varphi : X^3 \rightarrow [0, \infty)$ be a function and let $f : X \rightarrow Y$ be a mapping such that

$$\psi(x, y, z) := \sum_{j=0}^{\infty} \frac{1}{2^j} \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}, \frac{z}{2^j}\right) < \infty \quad (4.6)$$

and let $f : X \rightarrow Y$ be a mapping $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1\left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right)\right) \right\|_{\mathbb{Y}} \\ & \quad + \left\| \Gamma_2\left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z)\right) \right\|_{\mathbb{Y}} + \varphi(x, y, z) \quad (4.7) \end{aligned}$$

for all $x, y, z \in \mathbb{X}$.
Then there exists a unique additive mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbb{Y}} \leq \frac{1}{2(1 - |\Gamma_1|)} \psi(x, x, x) \quad (4.8)$$

for all $x \in \mathbb{X}$

Proof. Let $x = y = z$ in (4.7), we get

$$(1 - |\Gamma_1|) \|f(2x) - 2f(x)\| \leq \varphi(x, x, x) \quad (4.9)$$

for all $x \in X$. So

$$\left\| f(x) - \frac{1}{2}f(2x) \right\| \leq \frac{1}{(1 - |\Gamma_1|)} \varphi(x, y, z)$$

for all $x \in X$. Hence

$$\begin{aligned} & \left\| \frac{1}{2^l}f(2^l x) - \frac{1}{2^m}f(2^m x) \right\| \\ & \leq \sum_{j=l}^{m-1} \left\| \frac{1}{2^j}f(2^j x) - \frac{1}{2^{j+1}}f(2^{j+1} x) \right\| \\ & \leq \sum_{j=l}^{m-1} \frac{1}{2^{j+1}(1 - |\Gamma_1|)} \varphi(2^j x, 2^j x, 2^j x) \end{aligned} \quad (4.10)$$

for all nonnegative integers m and l with $m > l$ and all $x \in \mathbb{X}$. It follows from (4.10) that the sequence $\left\{ \frac{1}{2^n}f(2^n x) \right\}$ is a Cauchy sequence for all $x \in \mathbb{X}$. Since \mathbb{Y} is complete, the sequence $\left\{ \frac{1}{2^n}f(2^n x) \right\}$ converges. So one can define the mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ by

$$H(x) := \lim_{n \rightarrow \infty} \frac{1}{2^n}f(2^n x)$$

Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (4.10), we get (4.8). The rest of the proof is similar to the proof of theorem 4.1.

From proving the theorems we have consequences:

Corollary 4.3. *Let $r > 1$ and θ be nonnegative real numbers and let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and*

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\ & + \left\| \Gamma_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \theta \left(\|x\|^r + \|y\|^r + \|z\|^r \right) \end{aligned} \quad (4.11)$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique additive mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbb{Y}} \leq \frac{3\theta}{(2^r - 2)(1 - |\Gamma_1|)} \varphi(x, x, x) \quad (4.12)$$

for all $x \in \mathbb{X}$

Corollary 4.4. Let $r < 1$ and θ be nonnegative real numbers and let $f : \mathbb{X} \rightarrow \mathbb{Y}$ be a mapping satisfy $f(0) = 0$ and

$$\begin{aligned} & \left\| f\left(\frac{x+y}{2} + z\right) - f\left(\frac{x+y}{2}\right) - f(z) \right\|_{\mathbb{Y}} \\ & \leq \left\| \Gamma_1 \left(f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+y}{2} - z\right) - 2f\left(\frac{x+y}{2}\right) \right) \right\|_{\mathbb{Y}} \\ & + \left\| \beta_2 \left(2f\left(\frac{x+y}{2^2} + \frac{z}{2}\right) - f\left(\frac{x+y}{2}\right) - f(z) \right) \right\|_{\mathbb{Y}} + \theta \left(\|x\|^r + \|y\|^r + \|z\|^r \right) \end{aligned} \quad (4.13)$$

for all $x, y, z \in \mathbb{X}$.

Then there exists a unique additive mapping $H : \mathbb{X} \rightarrow \mathbb{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbb{Y}} \leq \frac{3\theta}{(2^r - 2)(1 - |\Gamma_1|)} \varphi(x, x, x) \quad (4.14)$$

for all $x \in \mathbb{X}$

5. CONCLUSION

In this paper, I have shown that the solutions of the Γ_1, Γ_2 -functional inequalities are additive mappings. The Hyers-Ulam stability for these given from theorems. These are the main results of the paper, which are the generalization of the results [3],[4],[14],[21].

REFERENCES

- [1]. T. Aoki, On the stability of the linear transformation in Banach space, J. Math. Soc. Japan 2(1950), 64-66...
- [2]. A.Bahyrycz, M. Piszczek, Hyers stability of the Jensen function equation, Acta Math. Hungar.,142 (2014),353-365.
- [3]. Ly V an An. Hyers-Ulam stability of functional inequalities with three variable in Banach spaces and Non-Archimedean Banach spaces International Journal of Mathematical Analysis Vol.13, 2019, no. 11. 519-53014), 296-310. <https://doi.org/10.12988/ijma.2019.9954>.
- [4]. Ly V an An. Hyers-Ulam stability of β -functional inequalities with three variable in non-Archimedean Banach spaces and complex Banach spaces International Journal of Mathematical Analysis Vol. 14, 2020, no. 5, 219 - 239 .HIKARI Ltd, www.m-hikari.com <https://doi.org/10.12988/ijma.2020.91169..><https://doi.org/10.12988/ijma.2019.9954>
- [5]. M.Balcerowski, On the functional equations related to a problem of z Boros and Z. Dro'czy, Acta Math. Hungar.,138 (2013), 329-340. .

- [6]. L. CADARIU ?, V. RADU, Fixed points and the stability of Jensens functional equation, J. Inequal. Pure Appl. Math. 4, no. 1, Art. ID 4 (2003). .
- [7]. J. DIAZ, B. MARGOLIS, A fixed point theorem of the alternative for contractions on a generalized complete metric space, Bull. Am. Math. Soc. 74 (1968), 305-309. .
- [8]. W Fechner, On some functional inequalities related to the logarithmic mean, Acta Math., Hungar., 128 (2010,)31-45, 303-309 . .
- [9]. W. Fechner, Stability of a functional inequalities associated with the Jordan-von Neumann functional equation, Aequationes Math. 71 (2006), 149-161
- [10]. Pascus. Gāvruta, A generalization of the Hyers-Ulam -Rassias stability of approximately ad- ditive mappings, Journal of mathematical Analysis and Aequations 184 (3) (1994), 431-436. <https://doi.org/10.1006/jmaa.1994.1211> .
- [11]. Attila Gila'nyi, On a problem by K. Nikodem, Math. Inequal. Appl., 5 (2002), 707-710.
- [12]. Attila Gila'nyi, Eine zur parallelogrammleichung äquivalente ungleichung, Aequations 5 (2002),707- 710. <https://doi.org/10.7153/mia-05-71> .
- [13]. Donald H. Hyers, On the stability of the functional equation, Proceedings of the National Academy of the United States of America, 27 (4) (1941), 222.<https://doi.org/10.1073/pnas.27.4.222>,
- [14]. Jung Rye Lee, Choonkil Park, and DongY unShin. Additive and quadratic functional in equalities in Non-Archimedean normed spaces, International Journal of Mathematical Analysis, 8 (2014), 1233- 1247.. <https://doi.org/10.12988/ijma.2019.9954>.
- [15]. L. Maligranda. Tosio Aoki (1910-1989). In International symposium on Banach and function spaces: 14/09/2006-17/09/2006, pages 1–23. Yokohama Publishers, 2008.
- [16]. D. MIHET, V. RADU, On the stability of the additive Cauchy functional equation in random normed spaces, J. Math. Anal. Appl. 343 (2008), 567-572.
- [17]. A. Najati and G. Z. Eskandani. Stability of a mixed additive and cubic functional equation in quasi-Banach spaces. J. Math. Anal. Appl., 342(2):1318–1331, 2008.
- [18]. C. Park,Y. Cho, M. Han. Functional inequalities associated with Jordan-von Newman-type additive functional equations , J. Inequality .Appl. ,2007(2007), Article ID 41820, 13 pages
- [19]. W. P and J. Schwaiger, A system of two in homogeneous linear functional equations, Acta Math. Hungar 140 (2013), 377-406 ..
- [20]. Jūrg Rātz On inequalities assosiated with the jordan-von neumann functional equation, Aequationes matheatiae, 66 (1) (2003), 191-200 <https://doi.org/10.1007/s00010-0032684-8>.
- [21]. Choonkil. Park. the stability an of additive (β_1, β_2) -functional inequality in Banach spaceJournal of mathematical Inequalities Volume 13, Number 1 (2019), 95-104. .
- [22]. Choonkil. Park. Additive β -functional inequalities, Journal of Nonlinear Science and Appl. 7(2014), 296-310. .
- [23]. Choonkil Park,. functional in equalities in Non-Archimedean normed spaces. Acta Mathematica Sinica, English Series, 31 (3), (2015), 353-366. <https://doi.org/10.1007/s10114-015-4278-5>

- [24]. C. PARK, Additive p -functional inequalities and equations, J. Math. Inequal. 9 (2015), 17-26. .
- [25]. C. PARK, Additive p -functional inequalities in non-Archimedean normed spaces, J. Math. Inequal. 9 (2015), 397-407 .
- [26]. Themistocles M. Rassias, On the stability of the linear mapping in Banach space, proceedings of the American Mathematical Society, 27 (1978), 297-300. <https://doi.org/10.2307/s00010-003-2684-8> .
- [27]. F. SKOF, Propriet locali e approssimazione di operatori, Rend. Sem. Mat. Fis. Milano 53 (1983),113-129. .
- [28]. S M. ULam. A collection of Mathematical problems, volume 8, Interscience Publishers. New York,1960.